

This is a repository copy of *Ultrafast polarization of an electron beam in an intense bichromatic laser field*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/156029/>

Version: Published Version

Article:

Seipt, Daniel, Del Sorbo, Dario, Ridgers, Christopher P. orcid.org/0000-0002-4078-0887 et al. (1 more author) (2019) Ultrafast polarization of an electron beam in an intense bichromatic laser field. *Physical Review A*. 061402. ISSN 1094-1622

<https://doi.org/10.1103/PhysRevA.100.061402>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Ultrafast polarization of an electron beam in an intense bichromatic laser field

Daniel Seipt^{1,*}, Dario Del Sorbo², Christopher P. Ridgers³ and Alec G. R. Thomas¹¹*The Gérard Mourou Center for Ultrafast Optical Science, University of Michigan, Ann Arbor, Michigan 48109, USA*²*High Energy Density Science Division, SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA*³*York Plasma Institute, Department of Physics, University of York, York YO10 5DD, United Kingdom*

(Received 26 April 2019; revised manuscript received 5 July 2019; published 4 December 2019)

Here, we demonstrate the radiative polarization of high-energy electron beams in collisions with ultrashort pulsed bichromatic laser fields. Employing a Boltzmann kinetic approach for the electron distribution allows us to simulate the beam polarization over a wide range of parameters and determine the optimum conditions for maximum radiative polarization. Those results are contrasted with a Monte Carlo algorithm where photon emission and associated spin effects are treated fully quantum mechanically using spin-dependent photon emission rates. The latter method includes realistic focusing laser fields, which allows us to simulate a near-term experimentally feasible scenario of an 8 GeV electron beam scattering from a 1 PW laser pulse and provide a measurement that would verify the ultrafast radiative polarization in high-intensity laser pulses that we predict. Aspects of spin-dependent radiation reaction are also discussed, with spin polarization leading to a measurable (5%) splitting of the energies of spin-up and spin-down electrons.

DOI: [10.1103/PhysRevA.100.061402](https://doi.org/10.1103/PhysRevA.100.061402)

One of the driving forces in the development of petawatt class laser systems [1–3] is related to novel laser-plasma based accelerator concepts. Recent laser wakefield acceleration experiments have demonstrated acceleration of electrons up to 8 GeV energy in a single stage [4] and may enable future novel TeV electron-positron colliders for high-energy physics [5,6]. Spin-polarized beams are crucial for high-energy collider applications, for instance in order to suppress the standard-model background in searches for new physics beyond the standard model [7,8]. Studying the dynamics of spin-polarized electrons in a plasma wakefield is therefore important [9–11]. Spin forces have been suggested to be important in both astrophysical systems and high-intensity laser-plasma interactions [12]. With increasing interest in high-intensity laser-plasma interactions, it is essential to understand lepton spin effects on the overall plasma dynamics through radiation reaction—because of the spin dependence in the photon emission rates—and electron-positron pair generation [13]. There has been notable recent interest in spin polarization of leptons in high-intensity laser interactions [14–16].

It is known that initially unpolarized lepton beams radiatively polarize in storage rings due to asymmetries in the rates for synchrotron emission, the so-called Sokolov-Ternov effect [17–19]. As a result, their projected spins end up being predominantly antialigned with the magnetic field direction. For most machines this is a slow process, with a timescale of minutes or hours, scaling as $T \propto \gamma/\chi^3$ [18,19], where $\chi = \sqrt{e^2 p \cdot F^2 \cdot p}/m^3 = \gamma|e|B/m^2 \ll 1$ is the quantum nonlinearity parameter, with the field strength tensor F and electron mass m , charge e , and four-momentum p^μ with Lorentz factor $\gamma = p^0/m$ [20].

We previously demonstrated that a similar spin polarization can occur for electrons circulating at the magnetic nodes of two colliding intense laser pulses [21], with polarization timescales of few femtoseconds for laser intensities exceeding $5 \times 10^{22} \text{ W/cm}^2$. However, these orbits are unstable in general [22]. By first accelerating electrons to high energy and then colliding them with a laser pulse, it is possible to achieve $\chi \gtrsim 1$ with current PW class high-intensity lasers operating at intensities $\sim 10^{21} \text{ W/cm}^2$ and study radiative spin polarization in the strongly quantum regime. This laser-electron-beam collider setup was used in seminal SLAC E-144 experiments demonstrating nonlinear Compton scattering [23] and electron-positron pair production [24] and has recently been used to observe quantum radiation reaction [25,26]. In Ref. [22], we discussed a polarization dependence of the radiation reaction force, related to the fact that spin-down electrons radiate more power than spin-up electrons.

In this Rapid Communication, we propose using bichromatic laser fields to polarize an electron beam and predict a measurable modification of the resulting quantum radiation reaction, Fig. 1. Spin-dependent radiation-reaction effects are described in our model using (i) a kinetic approach where we solve a Boltzmann equation for distribution functions of spin-polarized electrons and (ii) a quasiclassical particle tracking approach where electrons are pushed classically between photon emissions, and the emissions are treated fully quantum mechanically using a Monte Carlo algorithm employing spin-dependent photon emission rates. We determine optimum parameters for maximum radiative polarization, and discuss spin-dependent radiation reaction leading to a measurable splitting of the mean electron energies.

For multi-GeV electrons colliding with a high-power laser pulse the emission of gamma photons can be described as nonlinear Compton scattering using strong-field QED in the Furry picture [13,27]. The quantum-radiation dominated regime is

*dseipt@umich.edu




FIG. 1. Electrons propagating through a bichromatic laser pulse perform spin flips dominantly in certain phases φ of the field: Electrons initially polarized along the $+y$ direction (yellow trajectories) flip their spin to down (trajectory colored purple) dominantly when $B_y > 0$, and this is where 1ω and 2ω add constructively (blue contours). The opposite spin flip dominantly happens when $B_y < 0$ where the 1ω and 2ω components of the laser are out of phase (orange contours). Therefore χ is larger for $\uparrow\downarrow$ -flips and many more of those flips happen than the $\downarrow\uparrow$ -flips, causing a polarization of the beam.

reached for normalized laser amplitude $a_0 = eE/m\omega \gg 1$ and $\chi \sim 1$. The short photon formation length implies that quantum emissions can be described as incoherent events in a locally constant field approximation (LCFA). Hence, LCFA photon emission rates are customarily employed in simulations [28–36], but see also [37–40].

In a laser pulse, the magnetic field oscillates, and so in contrast with the case of a static magnetic field, the polarization built up in one half-cycle is mostly lost during the following half-cycle. In Ref. [41] we investigated an laser-electron-beam collider scenario, where an asymmetry between the two half-cycles is introduced by using ultrashort subcycle laser pulses. In that case, however, only a very small degree of polarization could be achieved due to the short interaction duration. A similar effect had been predicted due to electron self-interaction [42]. In Ref. [41] we also showed the agreement between the LCFA and full QED S -matrix calculations for spin-polarization effects. In the scheme proposed here, the addition of a 2ω component to the laser breaks the symmetry and allows for net radiative polarization of the beam without the necessity of subcycle pulses.

For electrons in an external field the canonical choice for the spin quantization axis is the spacelike axial four-vector field $\zeta = \zeta(x, p) = \tilde{f} \cdot p / (p \cdot \tilde{f}^2 \cdot p)^{1/2} = (\mathbf{u} \cdot \boldsymbol{\zeta}^{\text{RF}}, \boldsymbol{\zeta}^{\text{RF}} + \mathbf{u}(\mathbf{u} \cdot \boldsymbol{\zeta}^{\text{RF}})/[\gamma(1 + \gamma)])$ [19,43], where $\boldsymbol{\zeta}^{\text{RF}}$ points along the magnetic field in the rest frame of the electron, and with \tilde{f} the dual of the normalized field strength tensor $f = eF/m$. During the emission of a photon the electron spin will be projected onto a quantum eigenstate, with spin projection eigenvalue ± 1 along the local value of ζ [14,19]. Between emissions, the electrons are assumed to follow classical trajectories, with four-velocity $u = (\gamma, \mathbf{u}) = p/m$ governed by the Lorentz force equation, $du/d\tau = f \cdot u$ [spin-gradient (Stern-Gerlach) forces are negligible here [22,44]], and the precession of the spin vector expectation value S governed by the Thomas—Bargman-Michel-Telegdi (T-BMT) equation

[45], $dS/d\tau = \frac{g_e}{2} f \cdot S + \frac{g_e - 2}{2} (S \cdot f \cdot u) u$, with gyromagnetic ratio g_e , and with initial conditions $S = \pm \zeta$.

While the spin described above is a property of an individual electron, the polarization is a property of the whole beam [19], which can be defined as the ensemble average $\mathbb{S} = \langle S_y^{\text{RF}} \rangle \approx \langle S_y \rangle$ for the scattering scenario discussed in this Rapid Communication.

Assuming the bichromatic laser field is a plane wave propagating along the negative z axis and polarized along the x axis, then the magnetic field, $\boldsymbol{\zeta}^{\text{RF}}$, and ζ all have nonvanishing y components only. We can therefore define two distributions, n^s , $s = \pm 1$, of electrons with spins projected on the y axis. The evolution of the distributions n^s and transitions between spin states, $n^{+1} \leftrightarrow n^{-1}$, may be described by Boltzmann equations with transitions due to radiation emission described by means of a collisionlike integrodifferential operator [46,47].

For electrons colliding with a plane-wave laser pulse, the light-front momentum $p^+ = p^0 + p^z$ is conserved under the classical Lorentz equation, with their dynamics solely determined by quantum radiation reaction effects [48]. We can thus write the kinetic equations for the distribution functions, $n^s(p^+, \varphi)$, in the one-dimensional (1D) approximation as

$$\frac{\partial n^s}{\partial \varphi} = \sum_{s'} \left[\int_0^\infty dk^+ n^{s'}(q^+) \frac{d\mathbb{R}^{s's_\zeta}}{dk^+}(q^+) - n^s \mathbb{R}^{s's_\zeta} \right], \quad (1)$$

with laser phase $\varphi = \omega(t + z)$ and $q^+ = p^+ + k^+$. The spin-dependent probability per unit laser phase for photon emission, i.e., the LCFA photon emission rate, appearing in the collision operator in (1) is given by [41]

$$\begin{aligned} \frac{d\mathbb{R}^{s_\zeta s_{\zeta'}}}{dk^+} = & -\frac{\alpha m^2}{\omega(p^+)^2} \left[(1 + s_\zeta s_{\zeta'}) \text{Ai}_1(z) \right. \\ & + (g + s_\zeta s_{\zeta'}) \frac{2\text{Ai}'(z)}{z} \\ & \left. + \left(s_\zeta t + s_{\zeta'} \frac{t}{1-t} \right) \frac{\text{Ai}(z)}{\sqrt{z}} \right]. \end{aligned} \quad (2)$$

Here s_ζ ($s_{\zeta'}$) denotes the value of the electron spin projection along the magnetic field in the instantaneous electron rest frame before (after) the photon emission, e.g., $\zeta = \tilde{f} \cdot p / (p \cdot \tilde{f}^2 \cdot p)^{1/2}$ with p as the instantaneous electron four-momentum prior to photon emission, and $\chi_p = (p \cdot \tilde{f}^2 \cdot p)^{1/2} / m^4$. In Eq. (1) $s_\zeta^{(i)} = s^{(i)} \text{sgn}(B_y)$.

The argument of the Airy functions, its derivative and integral $\text{Ai}_1(z) = \int_z^\infty dx \text{Ai}(x)$ is $z = \{t/[\chi_p(1-t)]\}^{2/3}$, with the normalized light-front momentum of the emitted photon $t = k^+/p^+ \in (0, 1)$, and $g = 1 + t^2/(2-2t)$. For typical high-energy electron-beam laser collisions ζ agrees with the local second binormal Frenet-Serret vector up to terms $(\text{tr} \tilde{f} \cdot f)^2 / m^4 \chi^2 \ll 1$ ensuring that s_ζ is a constant of motion in a constant crossed field, which is required for the applicability of the spin-dependent LCFA rates (2) in simulation codes [41,49,50].

The 1D approximation holds when $p_\perp \ll p^+$, the transverse quiver amplitude of electrons in the field $a_0/\gamma\omega$ is much smaller than the laser spot size w_0 [29,46], and that ζ is not precessing under the T-BMT equation. These kinetic equations were validated for the case of a uniform static




FIG. 2. Kinetic equation solution for a 7.8 GeV electron beam [4] colliding with a laser with $a_0 = 10.8$, $c_2 = 0.7$, 109 fs FWHM duration, $\chi_0 = 1$. Evolution of n^{+1} , n^{-1} , and S as a function of laser phase from top to bottom. n^{+1} means the spin is aligned along the $+y$ axis. The shape of the magnetic field B_y is shown as a thin gray curve in the lower plot.

magnetic field by comparison with Sokolov-Ternov rate equations [18,51].

The bichromatic field used to solve Eq. (1) is characterized by the function $\psi(\varphi) = [\cos \varphi + c_2 \cos(2\varphi + \Delta\varphi)] \cos^2(\varphi/4N_L) \Theta(2\pi N_L - |\varphi|)$, where $\varphi = \omega(t + z)$ is the phase of the fundamental light, fundamental frequency $\omega = 1.55$ eV, the relative phase $\Delta\varphi = 0$ is chosen for maximum asymmetry, and N_L is the number of laser cycles. Note that by this definition, the fraction of the total pulse energy in the second harmonic is $c_2^2/(1 + c_2^2)$. The nonzero field components are $B_y = -E_x = E_0 \psi(\varphi)$, with $\chi_0 = ep_0^+ E_0/m^3 = (p_0^+ \omega/m^2) a_0$. The number of cycles and field strength are related to the pulse duration, $T_{\text{FWHM}}[\text{fs}] = 2.43 N_L \lambda[\mu\text{m}]$, and laser power $P[\text{TW}] \simeq 0.0429 a_0^2 (1 + c_2^2) (w_0/\lambda)^2$.

Solutions of the kinetic equation as a function of laser phase φ are shown in Fig. 2 for $n^{+1}(0) = n^{-1}(0)$. The distributions $n^{\pm 1}$ in (a) and (b) show the energy loss due to quantum radiation reaction effects, with the blue curves representing the mean values. The blue curve in (c) depicts the buildup of the polarization of the electrons, reaching a final value of -7.9% .

In Fig. 3 we show a parameter scan of the achievable degree of polarization as a function of χ_0 and c_2 . It shows that $\chi_0 > 0.5$ is required to acquire some significant polarization for $T = 161$ fs. For larger values of $\chi_0 > 1.5$ it saturates for $c_2 \simeq 0.7$ at about $S \approx -17\%$. Note that $S \approx 0$ for a one-color laser pulse ($c_2 = 0$) for all χ_0 . The survey plot has been calculated for $m\gamma = 5$ GeV electrons interacting with a 161 fs laser pulse. As long as the change of polarization per laser cycle is small, a scaling relation can be found: For fixed χ_0 and c_2 , the same polarization degree will be




FIG. 3. Achievable degree of electron polarization as a function of χ_0 and bichromaticity parameter c_2 . Calculations have been performed for 5 GeV electrons colliding with a 161 fs laser pulse, i.e., $a_0(\chi_0 = 1) = 16.5$.

achieved for constant values of T/p^+ , which has been verified numerically.

In order to describe the electron polarization beyond the 1D plane-wave approximation, we extended the accepted quasichlassical model of high-intensity laser matter interactions [27,28,30–34,36] to include the spin degree of freedom. In this model, electrons follow classical trajectories between quantum events, at which point emission is treated stochastically using a Monte Carlo algorithm employing the spin-dependent rates, Eq. (2). For the classical propagation, we solve the covariant Lorentz force and T-BMT equations using a fourth-order Runge-Kutta solver for u^μ and S^μ , respectively.

Our Monte Carlo algorithm generalizes that presented in [30,52], to explicitly include the spin of the electrons. It works as follows: (i) Assign a final optical depth τ_{em} to each particle at the beginning of the simulation and directly after each emission. (ii) At each time step after the classical push, project the spin four-vector onto the local spin quantization direction $s_\zeta = -S \cdot \zeta$ and reduce the remaining optical depth by $\sum_{s'} \mathbb{R}^{s_\zeta s'} \omega u^+ \Delta\tau$. (iii) A photon is emitted when optical depth reaches zero. Draw a random number $r_1 \in [0, 1]$ uniformly. If $r_1 \leq \mathbb{R}^{s_\zeta, s'=-1} / \sum_{s'} \mathbb{R}^{s_\zeta s'}$ the electron will go to a spin-down state, $s' = -1$, and $s' = +1$ (up) otherwise. (iv) Sample the normalized energy of the emitted photon from the distribution $d\mathbb{R}^{s_\zeta s'}/dt$ using inverse transform sampling. (v) Electron recoil is assigned via $\mathbf{p}' = (1 - t)\mathbf{p}$ [36] (see also [53]), and the new spin four-vector is determined by $S'_\mu = s'_\zeta \zeta'_\mu$ where the new electron momentum \mathbf{p}' is used for calculating ζ' . The Monte Carlo code has been verified against solutions of the kinetic equation, and predicts the same degree of spin polarization, e.g., $S = -7.9\%$, for the case of Fig. 2.

The Monte Carlo approach allows simulation of realistic three-dimensional (3D) scattering scenarios with focused beams. We describe the laser as the superposition of two Gaussian beams in the paraxial approximation for the 1ω and 2ω frequency components with the same focal spot size, $w_0 = 8 \mu\text{m}$, and pulse duration, $T = 109$ fs, $a_0 = 10.8$, and $c_2 = 0.7$, adding up to 0.75 PW laser power. The two components have different Rayleigh ranges and Gouy phase, which means the 1ω and 2ω components will only be in phase close to the focal plane. The phase shift at the 1ω Rayleigh range is




FIG. 4. Phase-space distributions of scattered electrons (a)–(d) show a correlation of electron momenta and their spin polarization: Low-energy electrons are stronger polarized. The final energy distributions for up and down electrons (e) show the effect of spin-dependent radiation reaction: Down electrons lost more energy than spin-up electrons on average. Lower right: Polarization of postselected electrons with $\gamma < \gamma_*$ (left axis) and their relative fraction (right axis).

$\Delta\varphi \simeq 0.32$. This sets a lower limit on the focal spot size in principle.

We use the Monte Carlo algorithm to simulate the collision of this laser with a 7.8 GeV electron beam with 5% energy spread, 0.1 mrad angular divergence, and $2.4 \mu\text{m}$ beam size, i.e., $\chi_0 = 1$ [4]. For scattering from a focused beam, each electron has $S_y \approx \pm 1$ and it is straightforward to define the two fractions of electrons with spin up and down by $S_y \geq 0$. The polarization degree of the beam is calculated as an ensemble average $\bar{S} = \langle S_y \rangle$.

For this 3D simulation, the final degree of polarization of the whole beam reaches $\bar{S} = -6.6 \pm 0.4\%$, which is slightly lower than the prediction for the 1D plane-wave case. The statistical uncertainty is due to the finite number of 3×10^5 simulated electrons. Figure 4 shows phase spaces p^+-p_x (a) and p^+-p_y (b) of the final electrons, as well as the corresponding differential polarization degrees (b),(d). For the latter, the degree of polarization is calculated independently for each bin with at least 20 electrons. These show that the polarization is not uniform over phase space.

Figure 4(e) shows the energy spectra for electrons with $S_y \geq 0$, showing that there is a 6.6% excess of down electrons, yielding the stated polarization degree. Figure 4 also shows that the mean energy of the down electrons (blue vertical line) is lower than the mean energy of the up electrons (red vertical line) with a relative difference of about 5.4%. The difference in the mean energy of up and down electrons shows that radiation reaction effects are spin dependent. For $\chi \sim 1$, spin-down electrons radiate about 30% more power than the up electrons [17].

For an estimate of the spin-dependent radiation reaction, we describe the electron dynamics by the leading term of the quantum corrected Landau-Lifshitz equation for each fraction separately [47,54,55], yielding $\frac{d\Delta}{d\varphi} \simeq \frac{4\alpha\omega}{3m} a^2(\varphi)[(g_d - g_u)\bar{\gamma}^2 - (g_d + g_u)\bar{\gamma}\Delta]$, to linear order in the energy splitting $\Delta = \gamma_u - \gamma_d$, with spin-dependent Gaunt factors g_s [22]. The

difference in emitted power $\propto g_d - g_u > 0$ is partially canceled by the second term in the square brackets, yielding an equilibrium solution $\Delta/\bar{\gamma} \simeq (g_d - g_u)/(g_d + g_u) \sim 0.1$ for $\chi \sim 1$.

The phase-space distributions in Figs. 4(a)–4(d) indicate that the observed degree of polarization can be increased significantly by postselecting some electrons [see Fig. 4(f)]. If only low-energy electrons are selected with $\gamma < 2800$ (40% of all electrons), for instance by using a magnetic spectrometer, then their polarization degree is increased to $\bar{S} = -20\%$. A similar yet less pronounced enhancement can be achieved by restricting $|p_y|$ to small values.

We have shown that it is possible to spin polarize high-energy electron beams using intense bichromatic laser pulses. We used both a kinetic approach with a linear Boltzmann equation as well as a quasiclassical Monte Carlo approach. A parameter scan revealed that a beam polarization of around 17% can be achieved for $c_2 \simeq 0.7$ and $\chi \gtrsim 2$. We found phase-space correlations of the electron polarization which allow one to apply phase-space cuts for increasing the measured polarization. We discuss spin-dependent radiation reaction leading to a 5% mean energy splitting between up and down electrons. In Ref. [15], the generation of positrons with polarization degrees of up to 60% was proposed by exploiting the spin asymmetry in the strong-field pair-production process.

The proposed electron-beam polarization could be realized experimentally with present-day technology in an all-optical setup: Multi-GeV electron beams (up to 8 GeV) have been demonstrated using laser wakefield accelerators [4]. Similarly, one could realize this experiment with a PW laser at a conventional accelerator facility, for instance at SLAC or XFEL [56]. The petawatt class laser pulse required needs to be frequency doubled, and the required technology for generating the 2ω light is second harmonic generation in a nonlinear crystal (having typically 50% energy efficiency). Growth of crystals large enough to be compatible with PW

lasers has been developed [57]. The polarization of the GeV electron beam could be measured using (nonlinear) Compton scattering [19,58,59], with [59] anticipating an accuracy of 0.3% from multi-GeV electrons in a single shot.

Note added. Recently, Ref. [15] appeared on the arXiv, the subject of which is similar to our own.

This work supported by the US Army Research Office Grant No. W911NF-16-1-0044 (D.S., A.G.R.T.), by the DOE, LDRD program at SLAC, under Contract No. DE-AC02-76SF00515 (D.D.S.), and by the UK Engineering and Physical Sciences Research Council Grant No. EP/M018156/1 (C.P.R.).

-
- [1] S. Gales, K. A. Tanaka, D. L. Balabanski, F. Negoita, D. Stutman, O. Tesileanu, C. A. Ur, D. Ursescu, I. Andrei, S. Ataman, M. O. Cernaianu, L. D'Alessi, I. Dancus, B. Diaconescu, N. Djourellov, D. Filipescu, P. Ghenuche, D. G. Ghita, C. Matei, K. Seto, M. Zeng, and N. V. Zamfir, The extreme light infrastructure–nuclear physics (ELI-NP) facility: New horizons in physics with 10 PW ultra-intense lasers and 20 MeV brilliant gamma beams, *Rep. Prog. Phys.* **81**, 094301 (2018).
 - [2] J. H. Sung, H. W. Lee, J. Y. Yoo, J. W. Yoon, C. W. Lee, J. M. Yang, Y. J. Son, Y. H. Jang, S. K. Lee, and C. H. Nam, 42 PW, 20 fs Ti:sapphire laser at 0.1 Hz, *Opt. Lett.* **42**, 2058 (2017).
 - [3] E. Cartlidge, The light fantastic, *Science* **359**, 382 (2018).
 - [4] A. J. Gonsalves, K. Nakamura, J. Daniels, C. Benedetti, C. Pieronek, T. C. H. de Raadt, S. Steinke, J. H. Bin, S. S. Bulanov, J. van Tilborg, C. G. R. Geddes, C. B. Schroeder, Cs. Tóth, E. Esarey, K. Swanson, L. Fan-Chiang, G. Bagdasarov, N. Bobrova, V. Gasilov, G. Korn, P. Sasorov, and W. P. Leemans, Petawatt Laser Guiding and Electron Beam Acceleration to 8 GeV in a Laser-Heated Capillary Discharge Waveguide, *Phys. Rev. Lett.* **122**, 084801 (2019).
 - [5] W. Leemans and E. Esarey, Laser-driven plasma-wave electron accelerators, *Phys. Today* **62**(3), 44 (2009).
 - [6] C. B. Schroeder, C. Benedetti, E. Esarey, and W. P. Leemans, Laser-plasma-based linear collider using hollow plasma channels, *Nucl. Instrum. Methods Phys. Res., Sect. A* **829**, 113 (2016), 2nd European Advanced Accelerator Concepts Workshop (EAAC 2015).
 - [7] G. Moortgat-Pick, T. Abe, G. Alexander, B. Ananthanarayan, A. A. Babich, V. Bharadwaj, D. Barber, A. Bartl, A. Brachmann, S. Chen, J. Clarke, J. E. Clendenin, J. Dainton, K. Desch, M. Diehl, B. Dobos, T. Dorland, H. K. Dreiner, H. Eberl, J. Ellis, K. Flöttmann, H. Fraas, F. Franco-Solova, F. Franke, A. Freitas, J. Goodson, J. Gray, A. Han, S. Heinemeyer, S. Hesselbach, T. Hirose, K. Hohenwarter-Sodek, A. Juste, J. Kalinowski, T. Kernreiter, O. Kittel, S. Kraml, U. Langenfeld, W. Majerotto, A. Martinez, H. U. Martyn, A. Mikhailichenko, C. Milstene, W. Menges, N. Meyners, K. Mönig, K. Moffeit, S. Moretti, O. Nachtmann, F. Nagel, T. Nakanishi, U. Nauenberg, H. Nowak, T. Omori, P. Osland, A. A. Pankov, N. Paver, R. Pitthan, R. Pöschl, W. Porod, J. Proulx, P. Richardson, S. Riemann, S. D. Rindani, T. G. Rizzo, A. Schällicke, P. Schüler, C. Schwanenberger, D. Scott, J. Sheppard, R. K. Singh, A. Sopczak, H. Spiesberger, A. Stahl, H. Steiner, A. Wagner, A. M. Weber, G. Weiglein, G. W. Wilson, M. Woods, P. Zerwas, J. Zhang, and F. Zomer, Polarized positrons and electrons at the linear collider, *Phys. Rep.* **460**, 131 (2008).
 - [8] A. Vauth and J. List, Beam polarization at the ILC: Physics case and realization, *Int. J. Mod. Phys.: Conf. Ser.* **40**, 1660003 (2016).
 - [9] J. Vieira, C.-K. Huang, W. B. Mori, and L. O. Silva, Polarized beam conditioning in plasma based acceleration, *Phys. Rev. ST Accel. Beams* **14**, 071303 (2011).
 - [10] M. Wen, M. Tamburini, and C. H. Keitel, Polarized Laser-Wakefield-Accelerated Kiloampere Electron Beams, *Phys. Rev. Lett.* **122**, 214801 (2019).
 - [11] Y. Wu, L. Ji, X. Geng, Q. Yu, N. Wang, B. Feng, Z. Guo, W. Wang, C. Qin, X. Yan, L. Zhang, J. Thomas, A. Hützen, M. Büscher, P. Rakitzis, A. Pukhov, B. Shen, and R. Li, Polarized electron-beam acceleration driven by vortex laser pulses, *New J. Phys.* **21**, 073052 (2019).
 - [12] S. M. Mahajan, F. A. Asenjo, and R. D. Hazeltine, Comparison of the electron-spin force and radiation reaction force, *Mon. Not. R. Astron. Soc.* **446**, 4112 (2015).
 - [13] V. I. Ritus, Quantum effects of the interaction of elementary particles with an intense electromagnetic field, *J. Sov. Laser Res.* **6**, 497 (1985).
 - [14] Y.-F. Li, R. Shaisultanov, K. Z. Hatsagortsyan, F. Wan, C. H. Keitel, and J.-X. Li, Ultrarelativistic Electron-Beam Polarization in Single-Shot Interaction with an Ultraintense Laser Pulse, *Phys. Rev. Lett.* **122**, 154801 (2019).
 - [15] Y.-Y. Chen, P.-L. He, R. Shaisultanov, K. Z. Hatsagortsyan, and C. H. Keitel, Polarized positron beams via intense two-color laser pulses, *arXiv:1904.04110*.
 - [16] F. Wan, R. Shaisultanov, Y.-F. Li, K. Z. Hatsagortsyan, C. H. Keitel, and J.-X. Li, Ultrarelativistic polarized positron jets via collision of electron and ultraintense laser beams, *arXiv:1904.04305*.
 - [17] A. A. Sokolov and I. M. Ternov, *Synchrotron Radiation* (Akademie Verlag, Berlin, 1968).
 - [18] I. M. Ternov, Synchrotron radiation, *Phys. Usp.* **38**, 409 (1995).
 - [19] S. R. Mane, Yu M. Shatunov, and K. Yokoya, Spin-polarized charged particle beams in high-energy accelerators, *Rep. Prog. Phys.* **68**, 1997 (2005).
 - [20] We use natural Heaviside-Lorentz units with $\hbar = c = \epsilon_0 = 1$, with fine structure constant $\alpha = e^2/4\pi$. Lorentz indices are omitted whenever possible, i.e., $u^\mu \rightarrow u$, $f^{\mu\nu} \rightarrow f$, and we denote scalar products and contractions of tensor indices as $f^{\mu\nu}u_\nu \rightarrow f \cdot u$.
 - [21] D. Del Sorbo, D. Seipt, T. G. Blackburn, A. G. R. Thomas, C. D. Murphy, J. G. Kirk, and C. P. Ridgers, Spin polarization of electrons by ultraintense lasers, *Phys. Rev. A* **96**, 043407 (2017).
 - [22] D. Del Sorbo, D. Seipt, A. G. R. Thomas, and C. P. Ridgers, Electron spin polarization in realistic trajectories around the

- magnetic node of two counter-propagating, circularly polarized, ultra-intense lasers, *Plasma Phys. Controlled Fusion* **60**, 064003 (2018).
- [23] C. Bula, K. T. McDonald, E. J. Prebys, C. Bamber, S. Boege, T. Kotseroglou, A. C. Melissinos, D. D. Meyerhofer, W. Ragg, D. L. Burke, R. C. Field, G. Horton-Smith, A. C. Odian, J. E. Spencer, D. Walz, S. C. Berridge, W. M. Bugg, K. Shmakov, and A. W. Weidemann, Observation of Nonlinear Effects in Compton Scattering, *Phys. Rev. Lett.* **76**, 3116 (1996).
- [24] D. L. Burke, R. C. Field, G. Horton-Smith, J. E. Spencer, D. Walz, S. C. Berridge, W. M. Bugg, K. Shmakov, A. W. Weidemann, C. Bula, K. T. McDonald, E. J. Prebys, C. Bamber, S. J. Boege, T. Koffas, T. Kotseroglou, A. C. Melissinos, D. D. Meyerhofer, D. A. Reis, and W. Ragg, Positron Production in Multiphoton Light-By-Light Scattering, *Phys. Rev. Lett.* **79**, 1626 (1997).
- [25] J. M. Cole, K. T. Behm, E. Gerstmayr, T. G. Blackburn, J. C. Wood, C. D. Baird, M. J. Duff, C. Harvey, A. Ilderton, A. S. Joglekar, K. Krushelnick, S. Kuschel, M. Marklund, P. McKenna, C. D. Murphy, K. Poder, C. P. Ridgers, G. M. Samarin, G. Sarri, D. R. Symes, A. G. R. Thomas, J. Warwick, M. Zepf, Z. Najmudin, and S. P. D. Mangles, Experimental Evidence of Radiation Reaction in the Collision of a High-Intensity Laser Pulse with a Laser-Wakefield Accelerated Electron Beam, *Phys. Rev. X* **8**, 011020 (2018).
- [26] K. Poder, M. Tamburini, G. Sarri, A. Di Piazza, S. Kuschel, C. D. Baird, K. Behm, S. Bohlen, J. M. Cole, D. J. Corvan, M. Duff, E. Gerstmayr, C. H. Keitel, K. Krushelnick, S. P. D. Mangles, P. McKenna, C. D. Murphy, Z. Najmudin, C. P. Ridgers, G. M. Samarin, D. R. Symes, A. G. R. Thomas, J. Warwick, and M. Zepf, Experimental Signatures of the Quantum Nature of Radiation Reaction in the Field of an Ultraintense Laser, *Phys. Rev. X* **8**, 031004 (2018).
- [27] A. Di Piazza, C. Müller, K. Z. Hatsagortsyan, and C. H. Keitel, Extremely high-intensity laser interactions with fundamental quantum systems, *Rev. Mod. Phys.* **84**, 1177 (2012).
- [28] E. N. Nerush, I. Y. Kostyukov, A. M. Fedotov, N. B. Narozhny, N. V. Elkina, and H. Ruhl, Laser Field Absorption in Self-Generated Electron-Positron Pair Plasma, *Phys. Rev. Lett.* **106**, 035001 (2011).
- [29] S. S. Bulanov, C. B. Schroeder, E. Esarey, and W. P. Leemans, Electromagnetic cascade in high-energy electron, positron, and photon interactions with intense laser pulses, *Phys. Rev. A* **87**, 062110 (2013).
- [30] C. P. Ridgers, J. G. Kirk, R. Duclous, T. G. Blackburn, C. S. Brady, K. Bennett, T. D. Arber, and A. R. Bell, Modelling gamma-ray photon emission and pair production in high-intensity laser-matter interactions, *J. Comput. Phys.* **260**, 273 (2014).
- [31] M. Vranic, T. Grismayer, R. A. Fonseca, and L. O. Silva, Quantum radiation reaction in head-on laser-electron beam interaction, *New J. Phys.* **18**, 073035 (2016).
- [32] C. N. Harvey, A. Ilderton, and B. King, Testing numerical implementations of strong-field electrodynamics, *Phys. Rev. A* **91**, 013822 (2015).
- [33] A. Gonoskov, S. Bastrakov, E. Efimenko, A. Ilderton, M. Marklund, I. Meyerov, A. Muraviev, A. Sergeev, I. Surmin, and E. Wallin, Extended particle-in-cell schemes for physics in ultrastrong laser fields: Review and developments, *Phys. Rev. E* **92**, 023305 (2015).
- [34] B. King, N. Elkina, and H. Ruhl, Photon polarization in electron-seeded pair-creation cascades, *Phys. Rev. A* **87**, 042117 (2013).
- [35] B. King, Double Compton scattering in a constant crossed field, *Phys. Rev. A* **91**, 033415 (2015).
- [36] T. G. Blackburn, D. Seipt, S. S. Bulanov, and M. Marklund, Benchmarking semiclassical approaches to strong-field QED: Nonlinear Compton scattering in intense laser pulses, *Phys. Plasmas* **25**, 083108 (2018).
- [37] A. Di Piazza, M. Tamburini, S. Meuren, and C. H. Keitel, Implementing nonlinear Compton scattering beyond the local-constant-field approximation, *Phys. Rev. A* **98**, 012134 (2018).
- [38] A. Ilderton, B. King, and D. Seipt, Extended locally constant field approximation for nonlinear Compton scattering, *Phys. Rev. A* **99**, 042121 (2019).
- [39] A. Ilderton, A note on the conjectured breakdown of QED perturbation theory in strong fields, *Phys. Rev. D* **99**, 085002 (2019).
- [40] T. Podszus and A. Di Piazza, High-energy behavior of strong-field QED in an intense plane wave, *Phys. Rev. D* **99**, 076004 (2019).
- [41] D. Seipt, D. Del Sorbo, C. P. Ridgers, and A. G. R. Thomas, Theory of radiative electron polarization in strong laser fields, *Phys. Rev. A* **98**, 023417 (2018).
- [42] S. Meuren and A. Di Piazza, Quantum Electron Self-Interaction in a Strong Laser Field, *Phys. Rev. Lett.* **107**, 260401 (2011).
- [43] W. B. Berestetzki, E. M. Lifschitz, and L. P. Pitajewski, *Quantum Electrodynamics*, 2nd ed. (Pergamon, Oxford, 1982).
- [44] M. Tamburini, F. Pegoraro, A. Di Piazza, C. H. Keitel, and A. Macchi, Radiation reaction effects on radiation pressure acceleration, *New J. Phys.* **12**, 123005 (2010).
- [45] V. Bargmann, L. Michel, and V. L. Telegdi, Precession of the Polarization of Particles Moving in a Homogeneous Electromagnetic Field, *Phys. Rev. Lett.* **2**, 435 (1959).
- [46] N. Neitz and A. Di Piazza, Stochasticity Effects in Quantum Radiation Reaction, *Phys. Rev. Lett.* **111**, 054802 (2013).
- [47] C. P. Ridgers, T. G. Blackburn, D. Del Sorbo, L. E. Bradley, C. Slade-Lowther, C. D. Baird, S. P. D. Mangles, P. McKenna, M. Marklund, C. D. Murphy, and A. G. R. Thomas, Signatures of quantum effects on radiation reaction in laser-electron-beam collisions, *J. Plasma Phys.* **83**, 715830502 (2017).
- [48] C. Harvey, T. Heinzl, and M. Marklund, Symmetry breaking from radiation reaction in ultra-intense laser fields, *Phys. Rev. D* **84**, 116005 (2011).
- [49] E. Honig, E. L. Schucking, and C. V. Vishveshwara, Motion of charged particles in homogeneous electromagnetic fields, *J. Math. Phys.* **15**, 774 (1974).
- [50] D. Seipt and A. G. R. Thomas, A Frenet-Serret interpretation of particle dynamics in high-intensity laser fields, *Plasma Phys. Control. Fusion* **61**, 074005 (2019).
- [51] J. L. Synge and A. Schild, *Tensor Calculus* (Dover, New York, 1978).
- [52] R. Duclous, J. G. Kirk, and A. R. Bell, Monte Carlo calculations of pair production in high-intensity laser-plasma interactions, *Plasma Phys. Controlled Fusion* **53**, 015009 (2011).
- [53] T. G. Blackburn, D. Seipt, S. S. Bulanov, and M. Marklund, Radiation beaming in the quantum regime, [arXiv:1904.07745](https://arxiv.org/abs/1904.07745).

- [54] S. V. Bulanov, T. Z. Esirkepov, M. Kando, J. K. Koga, and S. S. Bulanov, Lorentz-Abraham-Dirac versus Landau-Lifshitz radiation friction force in the ultrarelativistic electron interaction with electromagnetic wave (exact solutions), *Phys. Rev. E* **84**, 056605 (2011).
- [55] A. G. R. Thomas, C. P. Ridgers, S. S. Bulanov, B. J. Griffin, and S. P. D. Mangles, Strong Radiation-Damping Effects in a Gamma-Ray Source Generated by the Interaction of a High-Intensity Laser with a Wakefield-Accelerated Electron Beam, *Phys. Rev. X* **2**, 041004 (2012).
- [56] The LUXE project at DESY, see for instance M. Wing's talk given at the workshop "Probing strong-field QED in electron-photon interactions," <https://indico.desy.de/indico/event/19493/> (2018).
- [57] A. E. Kokh, T. B. Bekker, K. A. Kokh, N. G. Kononova, V. A. Vlezko, P. Villeval, S. Durst, and D. Lupinski, Nonlinear LBO and BBO crystals for extreme light sources, in *2010 International Conference on Advanced Optoelectronics and Lasers* (IEEE, Piscataway, 2010) pp. 127–129.
- [58] A. Narayan, D. Jones, J. C. Cornejo, M. M. Dalton, W. Deconinck, D. Dutta, D. Gaskell, J. W. Martin, K. D. Paschke, V. Tvaskis, A. Asaturyan, J. Benesch, G. Cates, B. S. Cavness, L. A. Dillon-Townes, G. Hays, E. Ihloff, R. Jones, P. M. King, S. Kowalski, L. Kurchaninov, L. Lee, A. McCreary, M. McDonald, A. Micherdzinska, A. Mkrtchyan, H. Mkrtchyan, V. Nelyubin, S. Page, W. D. Ramsay, P. Solvignon, D. Storey, A. Tobias, E. Urban, C. Vidal, B. Waidyawansa, P. Wang, and S. Zhamkotchyan, Precision Electron-Beam Polarimetry at 1 GeV Using Diamond Microstrip Detectors, *Phys. Rev. X* **6**, 011013 (2016).
- [59] Y.-F. Li, R.-T. Guo, R. Shaisultanov, K. Z. Hatsagortsyan, and J.-X. Li, Electron Polarimetry with Nonlinear Compton Scattering, *Phys. Rev. Appl.* **12**, 014047 (2019).